

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

What is claimed is:

1. An isolated nucleic acid encoding a human MCH1 receptor or a mutant of such human MCH1 receptor which is activated by MCH or an analog or homolog thereof.
- 5 2. The nucleic acid of claim 1, wherein the nucleic acid is DNA.
3. The DNA of claim 2, wherein the DNA is cDNA.
- 10 4. The DNA of claim 2, wherein the DNA is genomic DNA.
5. The nucleic acid of claim 1, wherein the nucleic acid is RNA.
- 15 6. The nucleic acid of claim 1, wherein the human MCH1 receptor has an amino acid sequence identical to that encoded by the plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197).
- 20 7. The nucleic acid of claim 1, wherein the human MCH1 receptor comprises an amino acid sequence as shown in Figure 2 (SEQ ID NO: 2).
- 25 8. The nucleic acid of claim 1, wherein the mutant human MCH1 receptor comprises an amino acid sequence as shown in Figure 13 (SEQ ID NO: 27).

9. The nucleic acid of claim 1, wherein the mutant human MCH1 receptor comprises an amino acid sequence as shown in Figure 14 (SEQ ID NO: 28).
- 5 10. The nucleic acid of claim 1, wherein the mutant human MCH1 receptor comprises an amino acid sequence as shown in Figure 15 (SEQ ID NO: 29).
11. A purified human MCH1 receptor protein.
- 10 12. A vector comprising the nucleic acid of claim 1.
13. The vector of claim 12 adapted for expression in a cell which comprises the regulatory elements necessary for
15 expression of the nucleic acid in the cell operatively linked to the nucleic acid encoding the receptor so as to permit expression thereof, wherein the cell is a bacterial, amphibian, yeast, insect or mammalian cell.
- 20 14. The vector of claim 13, wherein the vector is a baculovirus.²
15. The vector of claim 12, wherein the vector is a plasmid.
- 25 16. The plasmid of claim 15 designated pEXJ.HR-TL231 (ATCC Accession No. 203197).
17. A cell comprising the vector of claim 13.

18. A cell of claim 17, wherein the cell is a non-mammalian cell.
- 5 19. A cell of claim 18, wherein the non-mammalian cell is a *Xenopus* oocyte cell or a *Xenopus* melanophore cell.
20. A cell of claim 17, wherein the cell is a mammalian cell.
- 10 21. A mammalian cell of claim 20, wherein the cell is a COS-7 cell, a 293 human embryonic kidney cell, a NIH-3T3 cell, a LM(tk-) cell, a mouse Y1 cell, or a CHO cell.
22. An insect cell comprising the vector of claim 13.
- 15 23. An insect cell of claim 22, wherein the insect cell is an Sf9 cell, an Sf21 cell or a *Trichoplusia ni* 5B-4 cell.
- 20 24. A membrane preparation isolated from the cell of claim 17.
- 25 25. A nucleic acid probe comprising at least 15 nucleotides which specifically hybridizes with a nucleic acid encoding a human MCH1 receptor, wherein the probe has a unique sequence corresponding to a sequence present within one of the two strands of the nucleic acid encoding a human MCH1 receptor present in plasmid pEXJ.HR-T231 (ATCC Accession No. 203197).

26. A nucleic acid probe comprising at least 15 nucleotides which specifically hybridizes with a nucleic acid encoding a human MCH1 receptor, wherein the probe has a unique sequence corresponding to a sequence present within (a) the nucleic acid sequence shown in Figure 1 (SEQ ID NO: 1) or (b) the reverse complement thereof.
27. The nucleic acid probe of claim 25 or 26, wherein the nucleic acid is DNA.
28. The nucleic acid probe of claim 25 or 26, wherein the nucleic acid is RNA.
29. An antisense oligonucleotide having a sequence capable of specifically hybridizing to the RNA of claim 5, so as to prevent translation of the RNA.
30. An antisense oligonucleotide having a sequence capable of specifically hybridizing to the genomic DNA of claim 4.
31. An antisense oligonucleotide of claim 29 or 30, wherein the oligonucleotide comprises chemically modified nucleotides or nucleotide analogues.
32. An antibody capable of binding to a human MCH1 receptor encoded by the nucleic acid of claim 1.
33. An agent capable of competitively inhibiting the binding of the antibody of claim 32 to a human MCH1 receptor.

34. An antibody of claim 32, wherein the antibody is a monoclonal antibody or antisera.

5 35. A pharmaceutical composition comprising (a) an amount of the oligonucleotide of claim 29 capable of passing through a cell membrane and effective to reduce expression of a human MCH1 receptor and (b) a pharmaceutically acceptable carrier capable of passing through the cell membrane.

10 36. A pharmaceutical composition of claim 35, wherein the oligonucleotide is coupled to a substance which inactivates mRNA.

15 37. A pharmaceutical composition of claim 36, wherein the substance which inactivates mRNA is a ribozyme.

20 38. A pharmaceutical composition of claim 35, wherein the pharmaceutically acceptable carrier comprises a structure which binds to a human MCH1 receptor on a cell capable of being taken up by the cells after binding to the structure.

25 39. A pharmaceutical composition of claim 35, wherein the pharmaceutically acceptable carrier is capable of binding to a human MCH1 receptor which is specific for a selected cell type.

30 40. A pharmaceutical composition which comprises an amount of the antibody of claim 32 effective to block binding of a ligand to a human MCH1 receptor and a

pharmaceutically acceptable carrier.

41. A transgenic, nonhuman mammal expressing DNA encoding a human MCH1 receptor of claim 1.

5

42. A transgenic, nonhuman mammal comprising a homologous recombination knockout of the native human MCH1 receptor.

10

43. A transgenic, nonhuman mammal whose genome comprises antisense DNA complementary to the DNA encoding a human MCH1 receptor of claim 1 so placed within the genome as to be transcribed into antisense mRNA which is complementary to mRNA encoding the human MCH1 receptor and which hybridizes to mRNA encoding the human MCH1 receptor, thereby reducing its translation.

15

44. The transgenic, nonhuman mammal of claim 41 or 42, wherein the DNA encoding the human MCH1 receptor additionally comprises an inducible promoter.

20

45. The transgenic, nonhuman mammal of claim 41 or 42, wherein the DNA encoding the human MCH1 receptor additionally comprises tissue specific regulatory elements.

25

46. A transgenic, nonhuman mammal of claim 41, 42, or 43, wherein the transgenic, nonhuman mammal is a mouse.

30

47. A process for identifying a chemical compound which

specifically binds to a mammalian MCH1 receptor which comprises contacting cells comprising DNA encoding, and expressing on their cell surface, the mammalian MCH1 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian MCH1 receptor, wherein the cells do not normally express the mammalian MCH1 receptor and the DNA encoding the mammalian MCH1 receptor (a) hybridizes to a nucleic acid having the defined sequence shown in Figure 1 (SEQ ID NO: 1) under low stringency conditions or a sequence complementary thereto and (b) is further characterized by its ability to cause a change in the pH of a culture of CHO cells when a MCH1 ligand is added to the culture and the CHO cells contain the nucleic acid which hybridized to the nucleic acid having the defined sequence or its complement.

48. A process for identifying a chemical compound which specifically binds to a mammalian MCH1 receptor which comprises contacting a membrane preparation from cells comprising DNA encoding, and expressing on their cell surface, the mammalian MCH1 receptor, with the compound under conditions suitable for binding, and detecting specific binding of the chemical compound to the mammalian MCH1 receptor, wherein the cells do not normally express the mammalian MCH1 receptor and the DNA encoding the mammalian MCH1 receptor (a) hybridizes to a nucleic acid having the defined sequence shown in Figure 1 (SEQ ID NO: 1) under low stringency conditions or a sequence complementary thereto and (b) is further characterized by its ability to cause a change in the pH of a culture of CHO cells when a MCH1 ligand is added to the culture and the CHO cells contain the nucleic acid which hybridized to the nucleic acid having the defined sequence or its complement.

49. The process of claim 47 or 48, wherein the mammalian MCH1 receptor is a human MCH1 receptor.

5 50. The process of claim 47 or 48, wherein the mammalian MCH1 receptor is a rat MCH1 receptor.

10 51. The process of claim 47 or 48, wherein the mammalian MCH1 receptor has substantially the same amino acid sequence as the sequence of the human MCH1 receptor encoded by plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197).

15 52. The process of claim 47 or 48, wherein the mammalian MCH1 receptor comprises substantially the same amino acid sequence as that shown in Figure 2 (SEQ ID NO: 2).

20 53. The process of claim 47 or 48, wherein the mammalian MCH1 receptor comprises the amino acid sequence shown in Figure 2 (SEQ ID NO: 2).

54. The process of claim 47 or 48, wherein the mammalian MCH1 receptor comprises the amino acid sequence shown in Figure 13 (SEQ ID NO: 27).

25 55. The process of claim 47 or 48, wherein the mammalian MCH1 receptor comprises the amino acid sequence shown in Figure 14 (SEQ ID NO: 28).

56. The process of claim 47 or 48, wherein the mammalian MCH1 receptor comprises the amino acid sequence shown in Figure 15 (SEQ ID NO: 29).

5 57. The process of claim 47 or 48, wherein the compound is not previously known to bind to a mammalian MCH1 receptor.

58. A compound identified by the process of claim 57.

10 59. A process of claim 47 or 48, wherein the cell is an insect cell.

15 60. The process of claim 47 or 48, wherein the cell is a mammalian cell.

61. The process of claim 60, wherein the cell is nonneuronal in origin.

20 62. The process of claim 61, wherein the nonneuronal cell is a COS-7 cell, 293 human embryonic kidney cell, a CHO cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell.

25 63. A process of claim 60, wherein the compound is a compound not previously known to bind to a mammalian MCH1 receptor.

64. A compound identified by the process of claim 63.

5 65. A process involving competitive binding for identifying
a chemical compound which specifically binds to a
mammalian MCH1 receptor which comprises contacting cells
expressing on their cell surface the mammalian MCH1
receptor, with both the chemical compound and a second
chemical compound known to bind to the receptor, and
10 separately with only the second chemical compound, under
conditions suitable for binding of both compounds, and
detecting specific binding of the chemical compound to
the mammalian MCH1 receptor, a decrease in the binding
of the second chemical compound to the mammalian MCH1
15 receptor in the presence of the chemical compound
indicating that the chemical compound binds to the
mammalian MCH1 receptor, wherein the cells do not
normally express the mammalian MCH1 receptor and the DNA
encoding the mammalian MCH1 receptor (a) hybridizes to
20 a nucleic acid having the defined sequence shown in
Figure 1 (SEQ ID NO: 1) under low stringency conditions
or a sequence complementary thereto and (b) is further
characterized by its ability to cause a change in the pH
of a culture of CHO cells when a MCH1 ligand is added to
the culture and the CHO cells contain the nucleic acid
25 which hybridized to the nucleic acid having the defined
sequence or its complement.

30 66. A process involving competitive binding for identifying
a chemical compound which specifically binds to a
mammalian MCH1 receptor which comprises contacting a
membrane preparation from cells expressing on their cell
surface the mammalian MCH1 receptor, with both the
chemical compound and a second chemical compound known
to bind to the receptor, and separately with only the
35 second chemical compound, under conditions suitable for
binding of both compounds, and detecting specific

binding of the chemical compound to the mammalian MCH1 receptor, a decrease in the binding of the second chemical compound to the mammalian MCH1 receptor in the presence of the chemical compound indicating that the chemical compound binds to the mammalian MCH1 receptor, wherein the cells do not normally express the mammalian MCH1 receptor and the DNA encoding the mammalian MCH1 receptor (a) hybridizes to a nucleic acid having the defined sequence shown in Figure 1 (SEQ ID NO: 1) under low stringency conditions or a sequence complementary thereto and (b) is further characterized by its ability to cause a change in the pH of a culture of CHO cells when a MCH1 ligand is added to the culture and the CHO cells contain the nucleic acid which hybridized to the nucleic acid having the defined sequence or its complement.

67. A process of claim 65 or 66, wherein the mammalian MCH1 receptor is a human MCH1 receptor or a mutant of such human MCH1 receptor which is activated by MCH or an analog or homolog thereof.

68. A process of claim 65 or 66, wherein the mammalian MCH1 receptor is a rat MCH1 receptor.

69. The process of claim 65 or 66, wherein the cell is an insect cell.

70. The process of claim 65 or 66, wherein the cell is a mammalian cell.

71. The process of claim 70, wherein the cell is nonneuronal

in origin.

5 72. The process of claim 71, wherein the nonneuronal cell is a COS-7 cell, 293 human embryonic kidney cell, a CHO cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell.

73. The process of claim 70, wherein the compound is not previously known to bind to a mammalian MCH1 receptor.

10 74. A compound identified by the process of claim 73.

15 75. A method of screening a plurality of chemical compounds not known to bind to a mammalian MCH1 receptor to identify a compound which specifically binds to the mammalian MCH1 receptor, which comprises

20 (a) contacting cells transfected with and expressing DNA encoding the mammalian MCH1 receptor with the plurality of compounds not known to bind specifically to the mammalian MCH1 receptor, under conditions permitting binding of compounds known to bind the mammalian MCH1 receptor;

25 (b) determining whether the binding of a compound known to bind to the mammalian MCH1 receptor is reduced in the presence of the compounds within the plurality of compounds, relative to the binding of the compound in the absence of the
30 plurality of compounds; and if so

- (c) separately determining the binding to the mammalian MCH1 receptor of compounds included in the plurality of compounds, so as to thereby identify the compound which specifically binds to the mammalian MCH1 receptor.

5

76. A method of screening a plurality of chemical compounds not known to bind to a mammalian MCH1 receptor to identify a compound which specifically binds to the mammalian MCH1 receptor, which comprises

10

- (a) contacting a membrane preparation from cells transfected with and expressing DNA encoding the mammalian MCH1 receptor with the plurality of compounds not known to bind specifically to the mammalian MCH1 receptor under conditions permitting binding of compounds known to bind the mammalian MCH1 receptor;

15

- (b) determining whether the binding of a compound known to bind to the mammalian MCH1 receptor is reduced in the presence of the compounds within the plurality of compounds, relative to the binding of the compound in the absence of the plurality of compounds; and if so

20

25

- (c) separately determining the binding to the mammalian MCH1 receptor of compounds included in the plurality of compounds, so as to thereby identify the compound which specifically binds to the mammalian MCH1 receptor.

30

77. A method of claim 75 or 76, wherein the mammalian MCH1 receptor is a human MCH1 receptor or a mutant of such human MCH1 receptor which is activated by MCH or an analog or homolog thereof.

5

78. A method of claim 75 or 76, wherein the mammalian MCH1 receptor is a rat MCH1 receptor.

10

79. A method of claim 75 or 76, wherein the cell is a mammalian cell.

80. A method of claim 79, wherein the mammalian cell is non-neuronal in origin.

15

81. The method of claim 80, wherein the non-neuronal cell is a COS-7 cell, a 293 human embryonic kidney cell, a LM(tk-) cell, a CHO cell, a mouse Y1 cell, or an NIH-3T3 cell.

20

82. A method of detecting expression of a mammalian MCH1 receptor by detecting the presence of mRNA coding for the mammalian MCH1 receptor which comprises obtaining total mRNA from the cell and contacting the mRNA so obtained with the nucleic acid probe of any of claims 25, 26, 27, or 28 under hybridizing conditions, detecting the presence of mRNA hybridizing to the probe, and thereby detecting the expression of the mammalian MCH1 receptor by the cell.

25

30

83. A method of detecting the presence of a mammalian MCH1 receptor on the surface of a cell which comprises

contacting the cell with the antibody of claim 32 under conditions permitting binding of the antibody to the receptor, detecting the presence of the antibody bound to the cell, and thereby detecting the presence of the mammalian MCH1 receptor on the surface of the cell.

5

84. A method of determining the physiological effects of varying levels of activity of human MCH1 receptors which comprises producing a transgenic, nonhuman mammal of claim 44 whose levels of human MCH1 receptor activity are varied by use of an inducible promoter which regulates human MCH1 receptor expression.

10

85. A method of determining the physiological effects of varying levels of activity of human MCH1 receptors which comprises producing a panel of transgenic, nonhuman mammals of claim 44, each expressing a different amount of human MCH1 receptor.

15

86. A method for identifying an antagonist capable of alleviating an abnormality, wherein the abnormality is alleviated by decreasing the activity of a human MCH1 receptor comprising administering a compound to the transgenic, nonhuman mammal of claim 41, 44, 45, or 46, and determining whether the compound alleviates the physical and behavioral abnormalities displayed by the transgenic, nonhuman mammal as a result of overactivity of a human MCH1 receptor, the alleviation of the abnormality identifying the compound as an antagonist.

20

25

30

87. An antagonist identified by the method of claim 86.

88. A pharmaceutical composition comprising an antagonist of claim 87 and a pharmaceutically acceptable carrier.

5 89. A method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a human MCH1 receptor which comprises administering to the subject an effective amount of the pharmaceutical composition of claim 88, thereby treating the abnormality.

10 90. A method for identifying an agonist capable of alleviating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a human MCH1 receptor comprising administering a
15 compound to the transgenic, nonhuman mammal of claim 41, 44, 45, or 46, and determining whether the compound alleviates the physical and behavioral abnormalities displayed by the transgenic, nonhuman mammal, the alleviation of the abnormality identifying the compound
20 as an agonist.

91. An agonist identified by the method of claim 90.

25 92. A pharmaceutical composition comprising an agonist of claim 91 and a pharmaceutically acceptable carrier.

30 93. A method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a human MCH1 receptor which comprises administering to the subject an effective amount of the pharmaceutical composition of claim 92, thereby treating the abnormality.

94. A method for diagnosing a predisposition to a disorder associated with the activity of a specific mammalian allele which comprises:

5 (a) obtaining DNA of subjects suffering from the disorder;

(b) performing a restriction digest of the DNA with a panel of restriction enzymes;

10 (c) electrophoretically separating the resulting DNA fragments on a sizing gel;

15 (d) contacting the resulting gel with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MCH1 receptor and labeled with a detectable marker;

20 (e) detecting labeled bands which have hybridized to the DNA encoding a human MCH1 receptor of claim 1 labeled with a detectable marker to create a unique band pattern specific to the
25 DNA of subjects suffering from the disorder;

(f) preparing DNA obtained for diagnosis by steps (a) - (e); and

30 (g) comparing the unique band pattern specific to

the DNA of subjects suffering from the disorder from step (e) and the DNA obtained for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to the disorder if the patterns are the same.

5

95. The method of claim 94, wherein a disorder associated with the activity of a specific mammalian allele is diagnosed.

10

96. A method of preparing the purified human MCH1 receptor of claim 11 which comprises:

15

(a) inducing cells to express the human MCH1 receptor;

(b) recovering the human MCH1 receptor from the induced cells; and

20

(c) purifying the human MCH1 receptor so recovered.

97. A method of preparing the purified human MCH1 receptor of claim 11 which comprises:

25

(a) inserting nucleic acid encoding the human MCH1 receptor in a suitable vector;

(b) introducing the resulting vector in a suitable host cell;

5 (c) placing the resulting cell in suitable condition permitting the production of the isolated human MCH1 receptor;

(d) recovering the human MCH1 receptor produced by the resulting cell; and

10 (e) purifying the human MCH1 receptor so recovered.

15 98. A process for determining whether a chemical compound is a mammalian MCH1 receptor agonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian MCH1 receptor with the compound under conditions permitting the activation of the mammalian MCH1 receptor, and detecting an increase in mammalian MCH1 receptor activity, so as to thereby
20 determine whether the compound is a mammalian MCH1 receptor agonist.

25 99. A process for determining whether a chemical compound is a mammalian MCH1 receptor antagonist which comprises contacting cells transfected with and expressing DNA encoding the mammalian MCH1 receptor with the compound in the presence of a known mammalian MCH1 receptor agonist, under conditions permitting the activation of the mammalian MCH1 receptor, and detecting a decrease in
30 mammalian MCH1 receptor activity, so as to thereby determine whether the compound is a mammalian MCH1 receptor antagonist.

100. A process of claim 98 or 99, wherein the mammalian MCH1 receptor is a human MCH1 receptor or a mutant of such human MCH1 receptor which is activated by MCH or an analog or homolog thereof.

5

101. A process of claim 98 or 99, wherein the mammalian MCH1 receptor is a rat MCH1 receptor.

10

102. A pharmaceutical composition which comprises an amount of a mammalian MCH1 receptor agonist determined by the process of claim 98 effective to increase activity of a mammalian MCH1 receptor and a pharmaceutically acceptable carrier.

15

103. A pharmaceutical composition of claim 102, wherein the mammalian MCH1 receptor agonist is not previously known.

20

104. A pharmaceutical composition which comprises an amount of a mammalian MCH1 receptor antagonist determined by the process of claim 99 effective to reduce activity of a mammalian MCH1 receptor and a pharmaceutically acceptable carrier.

25

105. A pharmaceutical composition of claim 104, wherein the mammalian MCH1 receptor antagonist is not previously known.

30

106. A process for determining whether a chemical compound specifically binds to and activates a mammalian MCH1 receptor, which comprises contacting

cells producing a second messenger response and
expressing on their cell surface the mammalian MCH1
receptor, wherein such cells do not normally express
the mammalian MCH1 receptor, with the chemical
compound under conditions suitable for activation of
the mammalian MCH1 receptor, and measuring the
second messenger response in the presence and in the
absence of the chemical compound, a change in the
second messenger response in the presence of the
chemical compound indicating that the compound
activates the mammalian MCH1 receptor.

107. The process of claim 106, wherein the second
messenger response comprises chloride channel
activation and the change in second messenger is an
increase in the level of inward chloride current.

108. A process for determining whether a chemical
compound specifically binds to and inhibits
activation of a mammalian MCH1 receptor, which
comprises separately contacting cells producing a
second messenger response and expressing on their
cell surface the mammalian MCH1 receptor, wherein
such cells do not normally express the mammalian
MCH1 receptor, with both the chemical compound and
a second chemical compound known to activate the
mammalian MCH1 receptor, and with only the second
chemical compound, under conditions suitable for
activation of the mammalian MCH1 receptor, and
measuring the second messenger response in the
presence of only the second chemical compound and in
the presence of both the second chemical compound
and the chemical compound, a smaller change in the
second messenger response in the presence of both
the chemical compound and the second chemical
compound than in the presence of only the second

chemical compound indicating that the chemical compound inhibits activation of the mammalian MCH1 receptor.

- 5 109. The process of claim 108, wherein the second messenger response comprises chloride channel activation and the change in second messenger response is a smaller increase in the level of inward chloride current in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound.
- 10
- 15 110. A process of any of claims 106, 107, 108, or 109, wherein the mammalian MCH1 receptor is a human MCH1 receptor or a mutant of such human MCH1 receptor which is activated by MCH or an analog or homolog thereof.
- 20 111. A process of any of claims 106, 107, 108, or 109, wherein the mammalian MCH1 receptor is a rat MCH1 receptor.
- 25 112. The process of any of claims 106, 107, 108, 109, or 110, wherein the cell is an insect cell.
- 30 113. The process of any of claims 106, 107, 108, 109, or 110, wherein the cell is a mammalian cell.
114. The process of claim 113, wherein the mammalian cell is nonneuronal in origin.

115. The process of claim 114, wherein the nonneuronal cell is a COS-7 cell, CHO cell, 293 human embryonic kidney cell, NIH-3T3 cell or LM(tk-) cell.

5 116. The process of claim 106, 107, 108, or 109, wherein the compound is not previously known to bind to a mammalian MCH1 receptor.

117. A compound determined by the process of claim 116.

10 118. A pharmaceutical composition which comprises an amount of a mammalian MCH1 receptor agonist determined by the process of claim 106 or 107 effective to increase activity of a mammalian MCH1 receptor and a pharmaceutically acceptable carrier.

15 119. A pharmaceutical composition of claim 118, wherein the mammalian MCH1 receptor agonist is not previously known.

20 120. A pharmaceutical composition which comprises an amount of a mammalian MCH1 receptor antagonist determined by the process of claim 108 or 109 effective to reduce activity of a mammalian MCH1 receptor and a pharmaceutically acceptable carrier.

25 121. A pharmaceutical composition of claim 120, wherein the mammalian MCH1 receptor antagonist is not previously known.

122. A method of screening a plurality of chemical compounds not known to activate a mammalian MCH1 receptor to identify a compound which activates the mammalian MCH1 receptor which comprises:

5

(a) contacting cells transfected with and expressing the mammalian MCH1 receptor with the plurality of compounds not known to activate the mammalian MCH1 receptor, under conditions permitting activation of the mammalian MCH1 receptor;

10

(b) determining whether the activity of the mammalian MCH1 receptor is increased in the presence of the compounds; and if so

15

(c) separately determining whether the activation of the mammalian MCH1 receptor is increased by each compound included in the plurality of compounds, so as to thereby identify the compound which activates the mammalian MCH1 receptor.

20

123. A method of claim 122, wherein the mammalian MCH1 receptor is a human MCH1 receptor or a mutant of such human MCH1 receptor which is activated by MCH or an analog or homolog thereof.

25

124. A method of claim 122, wherein the mammalian MCH1 receptor is a rat MCH1 receptor.

30

125. A method of screening a plurality of chemical compounds not known to inhibit the activation of a mammalian MCH1 receptor to identify a compound which inhibits the activation of the mammalian MCH1 receptor, which comprises:

(a) contacting cells transfected with and expressing the mammalian MCH1 receptor with the plurality of compounds in the presence of a known mammalian MCH1 receptor agonist, under conditions permitting activation of the mammalian MCH1 receptor;

(b) determining whether the activation of the mammalian MCH1 receptor is reduced in the presence of the plurality of compounds, relative to the activation of the mammalian MCH1 receptor in the absence of the plurality of compounds; and if so

(c) separately determining the inhibition of activation of the mammalian MCH1 receptor for each compound included in the plurality of compounds, so as to thereby identify the compound which inhibits the activation of the mammalian MCH1 receptor.

126. A method of claim 125, wherein the mammalian MCH1 receptor is a human MCH1 receptor or a mutant of such human MCH1 receptor which is activated by MCH or an analog or homolog thereof.

127. A method of claim 125, wherein the mammalian MCH1 receptor is a rat MCH1 receptor.

5 128. A method of any of claims 123, 124, 125, 126, or 127, wherein the cell is a mammalian cell.

129. A method of claim 128, wherein the mammalian cell is non-neuronal in origin.

10 130. The method of claim 129, wherein the non-neuronal cell is a COS-7 cell, a 293 human embryonic kidney cell, a LM(tk-) cell or an NIH-3T3 cell.

15 131. A pharmaceutical composition comprising a compound identified by the method of claim 123 or 124 effective to increase mammalian MCH1 receptor activity and a pharmaceutically acceptable carrier.

20 132. A pharmaceutical composition comprising a compound identified by the method of claim 125 or 126 effective to decrease mammalian MCH1 receptor activity and a pharmaceutically acceptable carrier.

25 133. A method of treating an abnormality in a subject wherein the abnormality is alleviated by increasing the activity of a mammalian MCH1 receptor which comprises administering to the subject an amount of a compound which is a mammalian MCH1 receptor agonist effective to treat the abnormality.

134. A method of claim 133, wherein the abnormality is a regulation of a steroid or pituitary hormone disorder, an epinephrine release disorder, a gastrointestinal disorder, a cardiovascular disorder, an electrolyte balance disorder, hypertension, diabetes, a respiratory disorder, asthma, a reproductive function disorder, an immune disorder, an endocrine disorder, a musculoskeletal disorder, a neuroendocrine disorder, a cognitive disorder, a memory disorder, a sensory modulation and transmission disorder, a motor coordination disorder, a sensory integration disorder, a motor integration disorder, a dopaminergic function disorder, a sensory transmission disorder, an olfaction disorder, a sympathetic innervation disorder, pain, psychotic behavior, morphine tolerance, opiate addiction, an affective disorder, a stress-related disorder, a fluid-balance disorder, a seizure disorder, or migraine.

135. A method of treating an abnormality in a subject wherein the abnormality is alleviated by decreasing the activity of a mammalian MCH1 receptor which comprises administering to the subject an amount of a compound which is a mammalian MCH1 receptor antagonist effective to treat the abnormality.

136. A method of claim 135, wherein the abnormality is a regulation of a steroid or pituitary hormone disorder, an epinephrine release disorder, a gastrointestinal disorder, a cardiovascular disorder, an electrolyte balance disorder, hypertension, diabetes, a respiratory disorder, asthma, a reproductive function disorder, an immune disorder, an endocrine disorder, a musculoskeletal disorder, a neuroendocrine disorder, a cognitive

disorder, a memory disorder, a sensory modulation
and transmission disorder, a motor coordination
disorder, a sensory integration disorder, a motor
integration disorder, a dopaminergic function
disorder, a sensory transmission disorder, an
olfaction disorder, a sympathetic innervation
disorder, pain, psychotic behavior, morphine
tolerance, opiate addiction, an affective disorder,
a stress-related disorder, a fluid-balance disorder,
a seizure disorder, or migraine.

137. A process for making a composition of matter which
specifically binds to a mammalian MCH1 receptor
which comprises identifying a chemical compound
using the process of any of claims 47, 48, 65, 66,
75, or 76 and then synthesizing the chemical
compound or a novel structural and functional analog
or homolog thereof.

138. A process for making a composition of matter which
specifically binds to a mammalian MCH1 receptor
which comprises identifying a chemical compound
using the process of any of claims 98, 106, or 122
and then synthesizing the chemical compound or a
novel structural and functional analog or homolog
thereof.

139. A process for making a composition of matter which
specifically binds to a mammalian MCH1 receptor
which comprises identifying a chemical compound
using the process of any of claims 99, 108, or 125
and then synthesizing the chemical compound or a
novel structural and functional analog or homolog
thereof.

140. The process of any of claims 137, 138, or 139, wherein the mammalian MCH1 receptor is a human MCH1 receptor or a mutant of such human MCH1 receptor which is activated by MCH or an analog or homolog thereof.

141. The process of any of claims 137, 138, or 139, wherein the mammalian MCH1 receptor is a human MCH1 receptor.

142. A process for preparing a composition which comprises admixing a pharmaceutically acceptable carrier and a therapeutically effective amount of a chemical compound identified by the process of any of claims 47, 48, 65, 66, 75, or 76 or a novel structural and functional analog or homolog thereof.

143. A process for preparing a composition which comprises admixing a pharmaceutically acceptable carrier and a therapeutically effective amount of a chemical compound identified by the process of any of claims 98, 106, or 122 or a novel structural and functional analog or homolog thereof.

144. A process for preparing a composition which comprises admixing a pharmaceutically acceptable carrier and a therapeutically effective amount of a chemical compound identified by the process of any of claims 99, 108, or 125 or a novel structural and functional analog or homolog thereof.

145. The process of any of claims 142, 143, or 144,

wherein the mammalian MCH1 receptor is a human MCH1 receptor or a mutant of such human MCH1 receptor which is activated by MCH or an analog or homolog thereof.

5

146. The process of any of claims 142, 143, or 144, wherein the mammalian MCH1 receptor is a rat MCH1 receptor.

10

15

20

25

147. A process for determining whether a chemical compound is a human MCH1 receptor antagonist which comprises contacting cells transfected with and expressing DNA encoding the human MCH1 receptor with the compound in the presence of a known human MCH1 receptor agonist, under conditions permitting the activation of the human MCH1 receptor, and detecting a decrease in human MCH1 receptor activity, so as to thereby determine whether the compound is a human MCH1 receptor antagonist, wherein the DNA encoding the human MCH1 receptor comprises the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197), the known human MCH1 receptor agonist is MCH or a homolog or analog of MCH, and the cells do not express the MCH1 receptor prior to transfecting them.

30

148. A process for determining whether a chemical compound specifically binds to and inhibits activation of a human MCH1 receptor, which comprises separately contacting cells expressing on their cell surface the human MCH1 receptor and producing a second messenger response upon activation of the human MCH1 receptor, wherein such cells do not normally express the human MCH1 receptor and the DNA

encoding the human MCH1 receptor comprises the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197), with both the chemical compound and a second chemical compound known to activate the human MCH1 receptor, and with only the second chemical compound, under conditions suitable for activation of the human MCH1 receptor, and measuring the second messenger response in the presence of only the second chemical compound and in the presence of both the second chemical compound and the chemical compound, a smaller change in the second messenger response in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound indicating that the chemical compound inhibits activation of the human MCH1 receptor, wherein the second chemical compound is MCH or a homolog or analog of MCH.

149. The process of claim 148, wherein the second messenger response comprises chloride channel activation and the change in second messenger response is a smaller increase in the level of inward chloride current in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound.

150. A method of screening a plurality of chemical compounds not known to inhibit the activation of a human MCH1 receptor to identify a compound which inhibits the activation of the human MCH1 receptor, which comprises:

(a) contacting cells transfected with and expressing the human MCH1 receptor, wherein such cells do not normally express the human MCH1 receptor and the DNA encoding the human MCH1 receptor comprises the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197), with the plurality of compounds in the presence of a known human MCH1 receptor agonist, under conditions permitting activation of the human MCH1 receptor, wherein the known MCH1 receptor agonist is MCH or a homolog or analog of MCH;

(b) determining whether the activation of the human MCH1 receptor is reduced in the presence of the plurality of compounds, relative to the activation of the human MCH1 receptor in the absence of the plurality of compounds; and if so

(c) separately determining the extent of inhibition of activation of the human MCH1 receptor for each compound included in the plurality of compounds, so as to thereby identify the compound which inhibits the activation of the human MCH1 receptor.

151. The process of any of claims 147, 148 or 150, wherein the cell is an insect cell.

152. The process of any of claims 147, 148 or 150, wherein the cell is a mammalian cell.

153. The process of any of claims 147, 148 or 150, wherein the cell is a mammalian cell which is nonneuronal in origin.

5

154. The process of any of claims 147, 148 or 150, wherein the cell is a COS-7 cell, a CHO cell, a 293 human embryonic kidney cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell.

10

155. A process for making a composition of matter which specifically binds to a human MCH1 receptor which comprises identifying a chemical compound which specifically binds to the human MCH1 receptor and then synthesizing the chemical compound or a structural and functional analog or homolog thereof, wherein the chemical compound is identified as binding to the human MCH1 receptor by a process involving competitive binding which comprises contacting cells expressing on their cell surface the human MCH1 receptor, with both the chemical compound and a second chemical compound known to bind to the receptor, and separately with only the second chemical compound, under conditions suitable for binding of both compounds, and detecting the extent of specific binding of the chemical compound to the human MCH1 receptor, a decrease in the binding of the second chemical compound to the human MCH1 receptor in the presence of the chemical compound indicating that the chemical compound binds to the human MCH1 receptor, wherein the cells do not normally express the human MCH1 receptor, the human MCH1 receptor is encoded by nucleic acid comprising the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC Accession

15

20

25

30

35

No. 203197)), and the second chemical compound is MCH or a homolog or analog of MCH.

156. A process for making a composition of matter which
specifically binds to a human MCH1 receptor which
comprises identifying a chemical compound which
specifically binds to the human MCH1 receptor and
then synthesizing the chemical compound or a
structural and functional analog or homolog thereof,
wherein the chemical compound is identified as
binding to the human MCH1 receptor by a process
involving competitive binding which comprises
contacting a membrane preparation from cells
expressing on their cell surface the human MCH1
receptor, with both the chemical compound and a
second chemical compound known to bind to the
receptor, and separately with only the second
chemical compound, under conditions suitable for
binding of both compounds, and detecting the extent
of specific binding of the chemical compound to the
human MCH1 receptor, a decrease in the binding of
the second chemical compound to the human MCH1
receptor in the presence of the chemical compound
indicating that the chemical compound binds to the
human MCH1 receptor, wherein the cells do not
normally express the human MCH1 receptor, the human
MCH1 receptor is encoded by nucleic acid comprising
the sequence shown in Figure 1 (Seq. ID No. 1) or
contained in plasmid pEXJ.HR-TL231 (ATCC Accession
No. 203197), and the second chemical compound is MCH
or a homolog or analog of MCH.

157. A process for making a composition of matter which
is a human MCH1 receptor antagonist which comprises
identifying a chemical compound which is a human
MCH1 receptor antagonist and then synthesizing the

chemical compound or a structural and functional analog or homolog thereof, wherein the chemical compound is identified as a human MCH1 receptor antagonist by a process which comprises contacting cells transfected with and expressing DNA encoding the human MCH1 receptor with the compound in the presence of a known human MCH1 receptor agonist, under conditions permitting the activation of the human MCH1 receptor, and detecting a decrease in human MCH1 receptor activity, so as to thereby determine whether the compound is a human MCH1 receptor antagonist, wherein the cells do not normally express the human MCH1 receptor, the human MCH1 receptor is encoded by nucleic acid comprising the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197), and the known human MCH1 receptor agonist is MCH or a homolog or analog of MCH.

158. A process for making a composition of matter which specifically binds to and inhibits the activation of a human MCH1 receptor which comprises identifying a chemical compound which specifically binds to and inhibits the activation of the human MCH1 receptor and then synthesizing the chemical compound or a structural and functional analog or homolog thereof, wherein the chemical compound is identified as binding to and inhibiting the activation of the human MCH1 receptor by a process which comprises separately contacting cells expressing on their cell surface the human MCH1 receptor and producing a second messenger response upon activation of the human MCH1 receptor, wherein such cells do not normally express the human MCH1 receptor and the human MCH1 receptor is encoded by nucleic acid comprising the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC

Accession No. 203197), with both the chemical compound and a second chemical compound known to activate the human MCH1 receptor, and with only the second chemical compound, under conditions suitable for activation of the human MCH1 receptor, and measuring the second messenger response in the presence of only the second chemical compound and in the presence of both the second chemical compound and the chemical compound, a smaller change in the second messenger response in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound indicating that the chemical compound inhibits activation of the human MCH1 receptor, wherein the second chemical compound is MCH or a homolog or analog of MCH.

159. The process of claim 158, wherein the second messenger response comprises chloride channel activation and the change in second messenger response is a smaller increase in the level of inward chloride current in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound.

160. A process for preparing a composition which comprises identifying a chemical compound which specifically binds to a human MCH1 receptor, and then admixing a carrier and the chemical compound or a structural and functional analog or homolog thereof, wherein the chemical compound is identified as binding to the human MCH1 receptor by a process involving competitive binding which comprises contacting cells expressing on their cell surface the human MCH1 receptor, with both the chemical

compound and a second chemical compound known to bind to the receptor, and separately with only the second chemical compound, under conditions suitable for binding of both compounds, and detecting the extent of specific binding of the chemical compound to the human MCH1 receptor, a decrease in the binding of the second chemical compound to the human MCH1 receptor in the presence of the chemical compound indicating that the chemical compound binds to the human MCH1 receptor, wherein the cells do not normally express the human MCH1 receptor, the human MCH1 receptor is encoded by nucleic acid comprising the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197), and the second chemical compound is MCH or a homolog or analog of MCH.

161. A process for preparing a composition which comprises identifying a chemical compound which specifically binds to a human MCH1 receptor, and then admixing a carrier and the chemical compound or a structural and functional analog or homolog thereof, wherein the chemical compound is identified as binding to the human MCH1 receptor by a process involving competitive binding which comprises contacting a membrane preparation from cells expressing on their cell surface the human MCH1 receptor, with both the chemical compound and a second chemical compound known to bind to the receptor, and separately with only the second chemical compound, under conditions suitable for binding of both compounds, and detecting the extent of specific binding of the chemical compound to the human MCH1 receptor, a decrease in the binding of the second chemical compound to the human MCH1 receptor in the presence of the chemical compound indicating that the chemical compound binds to the

human MCH1 receptor, wherein the cells do not normally express the human MCH1 receptor, the human MCH1 receptor is encoded by nucleic acid comprising the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197), and the second chemical compound is MCH or a homolog or analog of MCH.

162. A process for preparing a composition which comprises identifying a chemical compound which is a human MCH1 receptor antagonist, and then admixing a carrier and the chemical compound or a structural and functional analog or homolog thereof, wherein the chemical compound is identified as a human MCH1 receptor antagonist by a process which comprises contacting cells transfected with and expressing DNA encoding the human MCH1 receptor with the compound in the presence of a known human MCH1 receptor agonist, under conditions permitting the activation of the human MCH1 receptor, and detecting a decrease in human MCH1 receptor activity, so as to thereby determine whether the compound is a human MCH1 receptor antagonist, wherein the cells do not normally express the human MCH1 receptor, the human MCH1 receptor is encoded by nucleic acid comprising the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197), and the known human MCH1 receptor agonist is MCH or a homolog or analog of MCH.

163. A process for preparing a composition which comprises identifying a chemical compound which specifically binds to and inhibits the activation of a human MCH1 receptor, and then admixing a carrier and the chemical compound or a structural and functional analog or homolog thereof, wherein the

chemical compound is identified as binding to and inhibiting activation of the human MCH1 receptor by a process which comprises separately contacting cells expressing on their cell surface the human MCH1 receptor and producing a second messenger response upon activation of the human MCH1 receptor, wherein such cells do not normally express the human MCH1 receptor and the human MCH1 receptor is encoded by nucleic acid comprising the sequence shown in Figure 1 (Seq. ID No. 1) or contained in plasmid pEXJ.HR-TL231 (ATCC Accession No. 203197), with both the chemical compound and a second chemical compound known to activate the human MCH1 receptor, and with only the second chemical compound, under conditions suitable for activation of the human MCH1 receptor, and measuring the second messenger response in the presence of only the second chemical compound and in the presence of both the second chemical compound and the chemical compound, a smaller change in the second messenger response in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound indicating that the chemical compound inhibits activation of the human MCH1 receptor, wherein the second chemical compound is MCH or a homolog or analog of MCH.

164. The process of claim 163, wherein the second messenger response comprises chloride channel activation and the change in second messenger response is a smaller increase in the level of inward chloride current in the presence of both the chemical compound and the second chemical compound than in the presence of only the second chemical compound.

165. The process of any of claims 155, 156, 157, 158, 160, 161, 162, or 163, wherein the cell is an insect cell.

5

166. The process of any of claims 155, 156, 157, 158, 160, 161, 162, or 163, wherein the cell is a mammalian cell.

10

167. The process of claim 166, wherein the mammalian cell is nonneuronal in origin.

15

168. The process of claim 167, wherein the nonneuronal cell is a COS-7 cell, a 293 human embryonic kidney cell, a CHO cell, a NIH-3T3 cell, a mouse Y1 cell, or a LM(tk-) cell.